

POSTERIOR TOTAL HIP ARTHROPLASTY

Dr. Brent Carlson
 Chippewa Valley Orthopedics & Sports Medicine
 1200 OakLeaf Way, Suite A 757 Lakeland Drive, Suite B
 Altoona, WI 54720 Chippewa Falls, WI 54729

Patient _____
 DOS _____

ACUTE CARE STAY	OUT-PATIENT THERAPY			NOTES:
<p>Week 0 Ankle Pumps, Quad Sets Gluteal Sets, Heel slides</p> <p>ROM restrictions: Greater than 90° Hip Flexion</p> <p>Bed mobility: May sleep on uninvolved side with pillow in between their knees.</p> <p>WBAT with assistive device. unless modified my MD. Progress gait as able, no limping.</p> <p>Stairs</p> <p>ADL Education: Use devices as needed for soft tissue discomfort needs. Pt should be encouraged to walk regularly.</p>	<p>1-3 weeks post-operative</p> <p>Continue post-op exercises</p> <p>AROM-supine, sliding with assist to encourage ROM within restrictions</p> <p>Long Arc Quad</p> <p>Hip adductor and abductor isometrics in hook lying</p> <p>Transverse abdominus isometrics and bridging</p> <p>Standing Hip Abduction, Extension and Flexion</p> <p>Bike, high seat</p> <p>Heel raises</p>	<p>4-6 weeks post-operative</p> <p>Continue stretches</p> <p>Continue strengthening</p> <p>Progress to: Hip Abduction with resistive tubing in hook lying.</p> <p>Clamshell</p> <p>Balance-double leg to single leg</p> <p>Total Gym</p> <p>Sub-max isotonic for hip, 1-5 pounds</p> <p>Walking activation</p> <ul style="list-style-type: none"> - March - Sidestep - Backwards <p>Bike</p> <p>Pool Therapy</p>	<p>7-12 weeks post-operative</p> <p>Progress ROM within restrictions and strength to WNL or equal to opposite extremity</p> <p>Progress strengthening of Quad and Hip groups</p> <p>Total gym with single leg</p> <p>Mini squats</p> <p>Wall sits</p> <p>Step-ups forward and lateral</p> <p>Balance</p> <p>D/C device when walking without a limp</p> <p>Address work and recreational functional activity demands.</p>	<p>These patients may be in a bit less pain than the anterior approach THA.</p> <p>There is no need for mandatory walker use due to less risk for iatrogenic fracture compared to anterior THA.</p> <p>Dr. Carlson does not detach the gluteus maximus and he makes certain these patients are stable through hyperflexion of knee to chest and internal rotation past 80° intraoperatively so hip precautions are meant to allow tissue relaxation post-op for 4 weeks.</p> <p>Progress to functional program as tolerated. Prepare for back to work, and recreational activities.</p> <p>This protocol should be interpreted as a continuum. If a patient is progressing ahead of the time schedules, advance them as tolerated.</p>
<p>Any Questions? Please contact: Northwoods Therapy Associates Altoona, WI Chippewa Falls, WI (715) 839-9266 (715) 723-5060</p>				